

OPTICAL PHENOMENA ASSOCIATED WITH THE LAUNCH OF A BALLISTIC MISSILE "BULAVA" DECEMBER 9, 2009

Yu.V. Platov ¹, S.A. Chernouss ²

1. Institute of Terrestrial Magnetism, Ionosphere and Radio Wave Propagation RAS, e-mail: yplatov@izmiran.ru
2. Polar Geophysical Institute KSC RAS, e-mail: chernouss@pgia.ru

An unexpected fairy show was occurred suddenly over the Norwegian Arctic December 9, 2009 early in the morning. A spiral glow larger than the Moon enchanted in the night sky above the polar provinces Tromsø, Finnmark, Trøndelag. Slowly rotating the spiral has become a significant part in the coverage of the sky light halo with disk-like shape. Blue-green beam rushing to the ground from the center of the halo, went on a rotation basis in the form of a corkscrew.

The lively discussion of the likely nature of the spectacle, observed people in Scandinavia was in this day on the network forums around. Norwegian media were overcrowded breathtaking images of eyewitnesses. The beautiful sight was observed residents throughout northern Norway, indicating its considerable height above the Earth.

Various hypotheses to explain the observed phenomenon were suggested. They extended from the invasion of aliens and God's signs to a more or less serious explanation. Thus, one of the hypotheses assumed that the "UFO over Norway has organized a new sky resort to attract attention or advertising the show was that the sky shone a spotlight. This trick is said to have been used previously, but now the organizers simply have to add capacity. According to another hypothesis, witty kids joked with a laser pointer on the streets.

The real explanation for the observed effects, as always, was much more prosaic.

ITAR-TASS in the press-service information and the Defense Ministry confirmed that on Dec. 9, 2009 at 06:45 UT (09:45 Moscow time) from a submarine TK-208 "Dmitry Donskoy", which was in the White Sea was made a test launch of a new Russian ballistic missile, sea-based Bulava from an underwater position. The test was unsuccessful. Following examination of the telemetry data revealed that the first two stages of the rocket worked in normal mode, but at a subsequent, third phase of the flight path there was a technical failure.

In the framework of bilateral agreements Russia has provided information on the technical characteristics of its newest missile the Bulava. It is three-stage rocket. Two first stages are solid-fuel ones, working on composite propellants. The third stage of the rocket, probably, also is solid-fuel one. Flaps for release of gases provide the maneuvers of the rocket.

Prior to the separation of the second stage of the rocket launch picture (Figure 1a) does not particularly different from the "classical" event. For comparison Fig. 1b shows a photograph of a missile launch «Minotaur» from Vandenberg on Sept. 23, 2005



Fig. 1. Rocket track visible as a result of the scattering of sunlight on the products of combustion.
a) during the White Sea launch b) during the Vanderberg launch

In either case, the field of flight the missiles were illuminated by sunlight, while observers were on the night side of Earth. The difference in light conditions is that the missile launch «Minotaur» took place at night, when observers were far east of the space rocket launch and the launch of the Bulava took place in the morning hours and, of course, it was impossible to watch it by witnesses, who reside in the regions to the west space of launch. It was not possible to observe this effect at the Russian territory (Murmansk region and Karelia) because at the time of the launch (~

9:45 MSK) the angle of the sun dipping below the horizon was small $\sim 6^\circ$, which corresponds to the height of the Earth's shadow about 30 km and at enough bright sky the launch was not seen.

A large amount of collected observations provided a unique opportunity for a correct assessment of geometrical parameters of the observed phenomena. This work was done in [1]. Figure 3 shows the obtained by the author estimates the size of various segments of the observed phenomena.




Fig 2. Composite picture of the development of optical phenomena observed in Norway during the launch the rocket on Jan. 9, 2009, obtained by combining a large number of photos.

The most interesting features of the optical effects that accompanied by the launch of the rocket, are presented: cyan (blue-green) the color of the missile track at altitudes up to ~ 100 km, the development of spiral structure in the region of the separation of the second stage of the rocket, the development of expanding around the location of the missiles "black hole" during the last stage of the phenomenon.

Development of the "black hole". The most obvious is the last of listed above effects. Actually there is no "black hole" is not developed. Engine stopped injected substances into the atmosphere just after turning off the third-stage and all the combustion products are ejected earlier, continued to fly apart. The rate of spread is determined by the speed of the combustion products into the atmosphere, their pressure and temperature. Engine injected substances into the atmosphere and injection velocity may be 2-3 km / sec [2] in normal mode of the engine in the upper atmosphere. While deviations from normal operation mode exhaust velocity should be considerably less than this value.

Development of a spiral. It is obvious that the spiral structure of gas-dust cloud formed during the third stage booster rocket, linked to its "somersaults", i.e. rotation is not only about its own axis, but also in the plane almost perpendicular to the line of sight. This is evidenced by the form of clouds, formed by the combustion products of the third stage. It is easy to find (Figure 3) that the compression ratio of the ellipse formed by the expanding spiral, i.e. ratio of its minor and major axes is ~ 0.85 . This corresponds to the angle between the plane of "tumbling" of the last stage and the direction of the line of sight from the place of observation $\sim 60^\circ$. A probable reason for such a "tumbling" may abort of stages separation.

Expansion of the combustion products is virtually a hemisphere [2, 3], when using of rocket engines in the upper atmosphere due to the high rate of underexpanded fuel (the ratio of pressure at the nozzle to the external pressure $> 10^5$). It is likely that the third stage engine took place in the off-design and products of incomplete combustion of fuel from the engine nozzle due to expire in the form of a dense jet with speeds significantly lesser design because, while expanding spiral and its branches enlarged in the transverse direction.




Fig. 3. Scheme of the spiral structure after the separation of the second stage.

Turquoise glow. Blue-green (turquoise) forms in the mesosphere frequently observed during solid-fuel rocket launches in the twilight conditions. Typically, these formations are sufficiently compact form "clouds", localized at the separation of rocket stages, and sometimes represent a characteristic diffuse winds "striated" trail - see Fig. 4a. Spectral observations conducted by auroral research programmes showed that the emissions of these formations are dominated by emissions of Al, AlO, Li (Fig. 4b).




Fig. 4. a) "Turquoise" smeared winds trail left by the passage of solid-fuel rocket, b) spectrum of gas-dust cloud formed by the combustion products of solid-propellant ballistic missile engine.

Metallic aluminum, lithium compounds and other components can be used in solid rocket fuels as admixtures to improve combustion stability [4, 5], and aluminum oxide can be formed by the oxidation of aluminum by atmospheric oxygen. It is also possible a different mechanism of formation of AlO. One of the main components of the combustion products of solid propellant missiles is aluminum oxide Al_2O_3 (~ 40% by weight [6]), which represents the fine fraction. At high temperatures (1300 K) as a result of accession to the molecules of aluminum oxide may be the formation of aluminum oxide AlO, located in a gaseous state. Blue-green fluorescence was also

observed repeatedly during active experiments with injection of high atmospheric trimethylaluminum or combustion of aluminum thermite charges [7, 8]. Mechanisms of chemiluminescent emission AIO not well understood, but it can be considered established that in the twilight conditions, the observed blue-green emission arise from resonant scattering of sunlight. In particular, this radiation is determined by the excitation of transitions $B^2 \Sigma \rightarrow ^2 \Sigma$ AIO molecules in the wavelength range 4374 Å - 5424 Å. [9]. Long-term "afterglow» AIO, ie glow for ~ an hour after the passage of the rocket is determined by the relatively small diffusion coefficient sufficiently heavy molecules.



Fig. 5. "Turquoise" track observed more than 40 minutes after the flight of the rocket.

Conclusion

The case study of rocket "Bulava" launch was discussed. Feature of this launch concerns to abort of stage separation. Other ones are the same as observed during solid propellant rocket launches. New model for formation of AIO cloud in the booster is proposed.

References

1. Spell T. «Estimation of the Location, Trajectory, Size, and Altitude of the "Norway Spiral" Phenomenon», M.Sc. Ocean Engineer (© SpellConsulting.com).
2. Molchanov A.G., Gas Dynamics of a Rocket Plume in the Upper Atmosphere, Proceeding 15th ESA Symposium on European Rocket and Balloon Programmes and Related Research, 28-31 May 2001, Biarritz, France, P. 375-379, 2001.
3. Simmons F.S. Rocket Exhaust Plume Phenomenology. California: Aerospace Press 2350 El Segundo, 2000.
4. Feodosyev V.I. Technical Fundamentals of rocket flight, Nauka, Moscow, 1981
5. Prisnyakov V.F., Dynamics of solid rocket motor fuel, Mashinostroenie, Moscow, 1984.
6. Environmental challenges and risks of exposure to rocket and space technology on the environment, reference manual, edited Adushkin VV, Kozlov SI and Petrova, AV, Ancile, M., 2000
7. Guseva N.N., Klyuyev O.F., calculation of the emission spectra of transition $A^2 \Sigma + \leftrightarrow X^2 \Sigma +$ AIO molecule to determine the temperature of the upper atmosphere, Proceedings of the Institute of Experimental Meteorology, Vol.6 (74), S. 52-58, 1977;
8. Gershenson Y.M., Grigorieva V.M., Maksyutov Sh.Sh., Chemistry artificial gas clouds in the Earth's upper atmosphere, Proceedings of the Institute of Experimental Meteorology, Vol. 21 (143), S. 3-31, 1990
9. Pierce R., Gaydon A., The identification of molecular spectra, Oxford, MA, 1949